

*Ibatova Sh.M.
Candidate of Medical Sciences,
Associate Professor, Samarkand State
Medical University,
Samarkand, Republic of Uzbekistan*

MODERN CONCEPTS OF THE ETIOLOGY AND RISK FACTORS FOR THE DEVELOPMENT OF PEPTIC ULCER DISEASE OF THE STOMACH AND DUODENUM

ANNOTATION

In recent years, there has been an increase in the incidence of duodenal ulcer and an increase in the number of complicated forms. Bleeding from duodenal ulcers is the most formidable complication and occupies one of the first places in the structure of mortality from duodenal ulcer. The anamnestic, clinical features and pH values in children with duodenal ulcer were studied. The study allows us to consider sex, age, heredity, erosive nature of gastroduodenitis, multiple ulcers, high bacterial contamination, low pH in the antroduodenal zone as factors for the complicated course of duodenal ulcer.

KEY WORDS: duodenal ulcer, factors, ulcerative bleeding, predictors of complications, children.

*Ибатова Ш.М.
Кандидат медицинских наук,
доцент Самаркандского государственного
медицинского университета,
Самарканд, Республика Узбекистан*

СОВРЕМЕННЫЕ ПРЕДСТАВЛЕНИЯ ОБ ЭТИОЛОГИИ И ФАКТОРАХ РИСКА РАЗВИТИЯ ЯЗВЕННОЙ БОЛЕЗНИ ЖЕЛУДКА И ДВЕНАДЦАТИПЕРСТНОЙ КИШКИ

АННОТАЦИЯ

В последние годы наблюдается увеличение заболеваемости язвой двенадцатиперстной кишки и числа осложненных форм. Кровотечение из язв двенадцатиперстной кишки является наиболее опасным осложнением и занимает одно из первых мест в структуре смертности от язвы двенадцатиперстной кишки. Были изучены анамнестические, клинические особенности и значения pH у детей с язвой двенадцатиперстной кишки. Исследование позволяет рассматривать пол, возраст, наследственность, эрозивный характер гастродуоденита, множественные язвы, высокую бактериальную контаминацию, низкий pH в антродуоденальной зоне как факторы осложненного течения язвы двенадцатиперстной кишки.

Ключевые слова: язва двенадцатиперстной кишки, факторы, язвенное кровотечение, предикторы осложнений, дети.

INTRODUCTION

In recent years, peptic ulcer due to the high prevalence, rejuvenation, aggravation of the clinical course of the disease and the lack of effectiveness of the therapy is an urgent problem of clinical medicine [1,4,5,9,15,18]. Among the etiological factors in the occurrence of peptic ulcer, heredity plays an important role. The role of hereditary burden is especially great in duodenal ulcer (DU), which is inherited in an autosomal dominant or autosomal recessive type, not sex-linked [3,4,5,10,12,14,19].

An important factor in the development of inflammatory and destructive changes is the specific infectious agent *Helicobacter pylori*. The frequency of infection of children with *Helicobacter pylori* infection is constantly increasing and reaches a maximum in the adult population, which makes it possible to consider *Helicobacteriosis* as the most common infection of mankind.

A feature of the course of peptic ulcer in modern conditions is a change in morphogenesis - an asymptomatic course or without pain is observed in more than 5% of patients; seasonal exacerbations are leveled, insufficient effectiveness or resistance to the therapy is noted, and complications of the disease often develop [5,11,17,20]. It is known that in 26-42% of patients with duodenal ulcer (DU) there are various complications, among which gastrointestinal bleeding accounts for 30-40% of cases, ulcer perforation - 21-27% [1,5,13,16].

Bleeding from duodenal ulcers is the most formidable complication and occupies one of the first places in the structure of mortality from duodenal ulcer. According to various authors, overall mortality is 10-14%, postoperative - from 12 to 35% [1,2,5-8].

The factors leading to the recurrence of the disease and the development of complications remain poorly understood, and the data published in the literature on the role of individual predictors in the development of complications are contradictory. The aspect of the problem of peptic ulcer associated with the search for predictors of the development of life-threatening complications is of undoubted relevance and is in demand in clinical practice.

PURPOSE OF THE STUDY

Study of the main causes of development of gastric ulcer and duodenal ulcer.

MATERIAL AND RESEARCH METHODS

In the 2nd clinic of the Samarkand Medical Institute, 38 sick children with duodenal ulcer (DU) in the acute phase were treated. The main group consisted of 12 (31.5%) patients with complicated course of the disease (bleeding) (PUD). The comparison group included 26 (68.4%) patients with uncomplicated DU (exacerbations of 1 or less during the year, no destructive complications ever). The study did not include patients with a history of taking ulcerogenic drugs (nonsteroidal anti-inflammatory drugs, glucocorticoids, etc.) during the last 3 months.

When distributing the examined by gender, the main group included 10 (83.3%) boys and 2 (16.7%) girls, in the comparison group, respectively, 18 (69%) and 8 (31%), respectively. The average age of patients at the time of the study was 16 ± 1.0 years in the group with complications, 12 ± 2.4 years without complications ($P < 0.01$).

Patients underwent: endoscopy, determination of the titer of IgG antibodies to HP in blood serum, intragastric pH-metry.

Endoscopy. Endoscopy was performed in all patients using the Olympus device (Japan) before treatment and after a course of antiulcer therapy.

Diagnosis of HP infection, determination of the presence of IgG antibodies to HP in the blood serum and their titer using the enzyme immunoassay "ImmunoComb HP IgG".

Intragastric pH monitoring was performed for all patients using the Gastroscan-AGM apparatus. At the same time, the average pH was evaluated. Patients with HP-positive DU received a standard 10-day eradication therapy regimen in the hospital: omeprazole 20 mg 2 times a day + amoxicillin 50 mg/kg 2 times a day + clarithromycin 15 mg/kg 2 times a day. Further treatment continued with Denol 120 mg 4 times a day for 3 weeks. Control EGDS was performed 21 days after the start of treatment until scarring of the ulcer. The effectiveness of eradication therapy was monitored using the respiratory Helic test 1 month after the end of treatment.

RESULTS AND ITS DISCUSSION

When studying the anamnesis of patients, it turned out that heredity for the underlying disease was aggravated. Thus, the presence of PU in close relatives of patients of the main group was detected in 7 (58.3%), and in 3 (25%) - a severe course, 4 (33.3%) of close relatives also had a history of bleeding. In sick children from the control group, hereditary burden was also traced 14 (46.7%), no complications were noted in relatives in this group.

It was established that the duration of the ulcer history in patients of the main group was 1.5 times less than in the comparison group, which indirectly indicates a more aggressive course of the disease in the former and, possibly, an asymptomatic debut.

In patients of the control group, the duration of the disease at the time of the examination was 3.1 ± 0.6 years, and in patients of the main group it was 1.4 ± 0.3 ($P < 0.01$). Relapses of the disease in the main group were observed less frequently than in the comparison group, 3 (25%) patients had destructive complications. According to the anamnestic study, there were no significant differences between the groups in terms of localization, irradiation of pain and its relationship with food intake. Night pains were more often recorded in the comparison group. Also, for the uncomplicated course of DU, moderate pains of typical localization were most characteristic.

The complicated course of DU was accompanied by less pronounced symptoms. The pain syndrome was quite typical in its localization, however, it had a lower intensity, the pains were more localized. With complicated DU, patients noted the absence of pain much more often than with DU (33.3% and 6.7%, respectively), less often - a significant intensity of the pain syndrome (25% and 40%). Seasonal exacerbations in the autumn-spring period were significantly more common in the group with PUF (56.7% and 33.3%, respectively).

Material and living conditions were unsatisfactory in 8 (66.7%) children with bleeding, and only in 12 (46%) of the comparison group. In endoscopic examination, antral gastritis and duodenitis were diagnosed in all patients. In 7 (58.3%) patients of the main group, an erosive nature of gastroduodenitis was noted.

Ulcerative defects in both groups of patients were localized mainly on the anterior and posterior walls of the duodenal bulb and much less frequently on the greater and lesser curvature. It should be noted that there were no statistically significant differences in the localization of

ulcerative defects in the duodenum in the studied groups. The average size of the ulcer in children with PUD was 3.1 ± 1.79 mm, which exceeds that in the control group (1.8 ± 1.14 mm) ($p > 0.05$).

When studying the acid-forming function of the stomach in children with DU, a state of hyperacidity was revealed in the area of the body of the stomach. The indicators of both the main and control groups significantly ($p < 0.01$; $p < 0.05$, respectively) differed from the standards. As for pH-metry in the antrum area, increased acid production (or a decrease in the alkalizing function of the stomach) is typical for children with a complicated course of the disease, in which this indicator was significantly ($p < 0.001$) lower than both the values of healthy and control groups.

According to the results of the study, HP infection was detected in 22 (84.6%) patients with PU and in all children of the main group. In patients with PU, a high titer of IgG to Hp was significantly more common (58.3% and 16.7%).

CONCLUSIONS

The following risk factors for the complicated course of duodenal ulcer in children were determined: male sex, senior school age, hereditary burden both for peptic ulcer and the presence of its complications, unsatisfactory material and living conditions, detection of erosive gastroduodenitis, large size of the ulcer and multiple ulcers. High titers of antibodies to *H. pylori*, low pH in children from the main group make it possible to include high bacterial contamination and hyperacidity in the group of predictors of complications of DU. Reducing the complications of the disease contributes to the clinical examination of patients in family clinics.

REFERENCES

1. Belonogov, N.I. Complications of peptic ulcer of the stomach and duodenum in urgent surgery /N.I. Belonogov // Materialy IX mezinarodnivedecko - praktickakonference "Moderni vymozhenostivedy- 2013". Lekafstvi: Praha. Publishing House "Education and Science" P.16-19.
2. Vidmanova T.A., Zhukova E.A., Viskova I.N. Efficiency of antihelicobacter therapy of duodenal ulcer in children. // Medical almanac.-2011.-№6.-C.182-184.
3. Komarchuk V.V. Evaluation of the acid-producing function of the stomach and diagnosis of gastroesophageal reflux in complicated forms of peptic ulcer. //J. Kharkiv surgical school.-2012.-№3(54)-p.50-52.
4. Midlenko, V.I. Algorithm for managing patients with ulcerative gastroduodenal bleeding // Kazan Medical Journal. - 2008. - No. 3. - P. 238-241.
5. Tsvetkova L.N., Goryacheva O.A., Nechaeva L.V. and others. The current course of peptic ulcer in children.// J. Pediatrics.-2007.-№6.-S.31-33.
6. Ibatova Sh. M., Mamatkulova F. Kh., Ruzikulov N.Y. The Clinical Picture of Acute Obstructive Bronchitis in Children and the Rationale for Immunomodulatory Therapy. International Journal of Current Research and Review. Vol 12 Issue 17. September 2020. - P.152-155.
7. Sh. M. Ibatova, F. Kh. Mamatkulova, N. B. Abdukadirova, Yu. A. Rakhmonov, M. M. Kodirova. Risk Factors for Development of Broncho-Ostructive Syndrome in Children. International Journal of Current Research and Review. Vol 12. Issue 23 December 2020. -P. 3-6.

8. Ibatova Sh.M., Mamatkulova F.Kh., Rakhmonov Y.A., Shukurova D.B., Kodirova M.M. Assessment of the Effectiveness of Treatment of Rachit in Children by Gas-Liquid Chromatography. International Journal of Current Research and Review. Vol 13, Issue 06, 20 March 2021. P.64-66.
9. Sh.M. Ibatova, F.Kh. Mamatkulova, D.S. Islamova. Efficiency of combined application of apricot oil and aevit as a regulator of lipase activity of blood serum in children with vitamin D-deficiency rickets. Journal of Critical Reviews. // ISSN- 2394-5125. VOL 7, ISSUE 11, 2020. P.1266-1274.
10. Mamatkulova F.Kh., Ibatova Sh.M., Baratova R.Sh., Ergashev A.Kh. State of immunity in chronic obstructive pulmonary disease in children. Asian Journal of Multidimensional Research (AJMR). Vol.10, Issue 3, March, 2021. P. 132-136.
11. Mamutova E.S. Ibatova Sh. M., Abdurasulov F.P., Some aspects of diagnostics of out-of-social pneumonia in children indications for hospitalization. EPRA International Journal of Research and Development (IJRD) Volume: 6 | Issue: 4 | April 2021. P. 242-244.
12. Ibatova Sh.M., Mamatkulova F.Kh., Mukhamadiev N.K. State of immunity in chronic obstructive pulmonary disease in children. Central Asian Journal of Medical and Natural Sciences Volume: 02 Issue: 05 | Sep-Oct 2021 ISSN: 2660-4159. P. 103-107.
13. Muhamadiev N.Q. Ibatova Sh.M., Efficiency of immunomodulating therapy in acute obstructive bronchitis in children. Central Asian Journal of Medical and Natural Sciences. Volume: 02 Issue: 02 | March-April 2021 ISSN: 2660-4159. P. 210-213.
14. E.S.Mamutova, Ibatova Sh.M., D.T. Rabbimova, N.B. Abdukadirova, M.M. Kadirova. Gas-chromatographic appraisal of application of apricot oil and aevit in complex therapy of vitamin D-deficiency rickets in children. International Scientific Journal Theoretical & Applied Science, 24.04.2019, Philadelphia, USA, P.333-336.
15. Ibatova Sh.M. Evaluation of the effectiveness of the use of apricot oil and aevit in the complex treatment of children with rickets // J. Bulletin of the Association of Physicians of Uzbekistan. - 2015. - No. 4. - P.50-53.
16. Ibatova Sh.M. Optimization of the treatment of rickets in children // J. Infection, immunity and pharmacology. - 2015. - No. 5. - S. 99-103.
17. D.S. Islamova, Ibatova Sh.M. Sh.O. Akhmedov. The use of cumin oil in the treatment of children with rickets // J. Problems of biology and medicine, Samarkand. - 2015. - No. 2 (83). - S. 50-52.
18. Sh.M. Ibatova, F.Kh. Mamatkulova, N.B. Abdukadirova, Kh.M. Oblokulov, F.A. Achilova. The effectiveness of the use of apricot oil in children with rickets. // Scientific and practical journal "Questions of Science and Education", Moscow, 2019, No. 27 (76), -P.40-46.
19. Sh.M. Ibatova, F.Kh. Mamatkulova Gas chromatographic evaluation of the use of apricot oil and aevit in the complex therapy of vitamin D-deficient rickets in children. Scientific journal "New day in medicine". No. 2 (30/2) 2020. -P. 265-267.
20. Ibatova Sh.M., F.Kh. Mamatkulova M.M. Kodirova. Evaluation of the effectiveness of treatment of rickets in children by gas-liquid chromatography Asian Journal of Multidimensional Research Vol. 9, Issue 10, October, 2020. P. 44-47.