

*Ibatova Sh..M.
Candidate of Medical Sciences,
Associate Professor Samarkand State
Medical University,
Samarkand, Republic of Uzbekistan*

*Ravshanova M.I.
second-year student, Faculty of Medicine,
Samarkand State Medical University,
Samarkand, Republic of Uzbekistan.*

*Shermatova E. Sh.
second -year student, Faculty of Medicine,
Samarkand State Medical University
Samarkand, Republic of Uzbekistan*

CLINICAL AND LABORATORY PROGNOSIS OF ACUTE GLOMERULONEPHRITIS IN CHILDREN WITH NEPHROTIC SYNDROME

ANNOTATION

In the development of chronic renal failure in children, the proportion of glomerulonephritis is still significant. Identification of chronic kidney pathology at the initial stage is important to improve the effectiveness of preventive and therapeutic measures aimed at slowing down the progression of the disease. We examined 76 patients with acute glomerulonephritis with nephrotic syndrome aged 2 to 5 years, including 31 boys and 45 girls. The development of the forecast algorithm was carried out by the method of sequential Wald analysis. A score of 10 was used as a threshold to conclude on the prognosis. The peculiarity of the course of glomerulonephritis in children was that the disease develops against the background of hereditary predisposition.

Key words: glomerulonephritis, nephrotic syndrome, patients, course, prognosis.

УДК 616.611-002

*Ибатова Ш.М.
Кандидат медицинских наук,
доцент Самаркандского государственного медицинского университета,
Самарканд, Республика Узбекистан*

*Равшанова М.И.
студентка второго курса медицинского факультета,
Самаркандский государственный медицинский университет,*

Самарканд, Республика Узбекистан.

Шерматова Э.Ш.

*студентка второго курса медицинского факультета,
Самаркандский государственный медицинский университет,
Самарканд, Республика Узбекистан*

КЛИНИЧЕСКИЙ И ЛАБОРАТОРНЫЙ ПРОГНОЗ ОСТРОГО ГЛОМЕРУЛОНЕФРИТА У ДЕТЕЙ С НЕФРОТИЧЕСКИМ СИНДРОМОМ

АННОТАЦИЯ

В развитии хронической почечной недостаточности у детей доля гломерулонефрита остается значительной. Выявление хронической патологии почек на ранней стадии важно для повышения эффективности профилактических и терапевтических мер, направленных на замедление прогрессирования заболевания. Мы обследовали 76 пациентов с острым гломерулонефритом с нефротическим синдромом в возрасте от 2 до 5 лет, в том числе 31 мальчика и 45 девочек. Разработка алгоритма прогнозирования проводилась методом последовательного анализа Вальда. В качестве порогового значения для заключения о прогнозе использовалась оценка в 10 баллов. Особенностью течения гломерулонефрита у детей является то, что заболевание развивается на фоне наследственной предрасположенности.

Ключевые слова: гломерулонефрит, нефротический синдром, пациенты, течение, прогноз.

INTRODUCTION

Glomerulonephritis is one of the main causes of chronic renal failure (CRF). The prevalence of the disease is growing in many countries of the world, and therefore it becomes extremely necessary to study the clinical course and prognosis of renal diseases [1,2,7,13,16].

In modern nephrology, age-related features of glomerular diseases are considered in two directions: differences between children and adults are studied, as well as features in children and adults depending on their age at the onset of the disease [6,7,9,12,14].

According to epidemiological studies, the structure of glomerular diseases may change. The development of nephrotic syndrome is associated with an increase in the permeability of the glomerular filtration barrier for macromolecules.

It is known that glomerular diseases can lead to a decrease in renal function. For a long time, impaired renal function can occur latently. Identification of chronic renal failure (CRF) in the initial stage is important to improve the effectiveness of preventive and therapeutic measures aimed at slowing down the progression of the disease [3,8,11,15,17]. Therefore, predicting the outcomes of glomerulonephritis in early childhood continues to be an urgent medical and social problem [5,10].

Purpose of the study: to determine the diagnostic significance of clinical and laboratory parameters in predicting acute glomerulonephritis with nephrotic syndrome in children.

Material and research methods. Patients with acute glomerulonephritis (76) with nephrotic syndrome aged 2 to 5 years were examined, including 30 boys and 44 girls. The patients were divided into 2 groups. The first group consisted of 35 patients who had no

recurrence of the disease for two or more years, the second group included 39 patients who did not achieve complete remission and developed a chronic form of the disease (CHN). The development of the forecast algorithm was carried out by the method of sequential Wald analysis [4]. A score of 10 was used as a threshold to conclude on the prognosis. The conclusion about the high probability of the transition of the disease to chronic glomerulonephritis was determined if the patient had more than 10 points. The development of the Wald sequential analysis algorithm was carried out by determining the frequency of symptoms in each group as a percentage. The evidence of statistical reliability of the difference in the frequency of symptoms is determined by the formula: $t = \frac{P1 - P2}{\sqrt{M221 - M22}}$. The proof of the independence of signs of detection of prognostic symptoms was the determination of the correlation coefficient for qualitative signs, which is calculated by the formula:

$H = \frac{ad - bc}{(a + b)(c + d)(a + c)(b + d)}$, where the letters indicate the signs. Correlation coefficient for quantitative traits, where "x" and "y" are average correlated traits. $r = \frac{Tx - Ty}{\sqrt{Tx^2 - Ty^2}}$. The determination of the relative probability and the calculation of the prognostic coefficient were carried out according to the formula: $PC = 10 \lg \frac{P1}{P2}$.

Indicators of unfavorable prognosis with a positive sign, because in the numerator, the frequency of symptoms characteristic of the group with developed CGN, and in the denominator, symptoms characteristic of the group with acute glomerulonephritis with long-term clinical and laboratory remission. Taking into account the hypothesis with a normal distribution, the information measure was calculated using the Kullback formula [2]. According to the degree of information content, the following signs were selected, listed in the table, where PC is a prognostic coefficient, $J(x_i)$ is information content.

Research results. A comparative assessment of hereditary burden revealed that a prognostically unfavorable sign of chronicity was the presence of kidney diseases in relatives in the pedigree, the prognostic coefficient (PC) was 9.9 and significantly informative +1.65. From the anamnestic data, there were frequent sore throats (PC = 15.3) with the highest information content among all other signs (6.12), food and drug allergies, parasitic infestations, PC = 5.2 and 6 with an information value of 1.35 and 6, respectively. 1.02. Thus, if the total score exceeds the threshold unit "10", the patient has an unfavorable prognosis and belongs to the high-risk group requiring appropriate therapy.

Of the laboratory studies, the most unfavorable prognostic indicator was partial renal dysfunction: hypo- and hyperkalemia, proteinuria more than 3 g/l per day. According to the coagulogram, there was plasma tolerance to heparin (7.6). According to the analysis of excretory urography, unfavorable factors of chronicity were an increase in the size of the kidneys (PC = 17), information content 1.76, an increase in the nephrographic effect (PC = 12) with information content 1.24. In the genesis of damage to renal structures in acute glomerulonephritis, a high information content of malondialdehyde was established with its increase from 6.8-8.4 nmol/mg/lipids in the cell membrane. Despite a slight difference in the indices of phosphatidylcholine, phosphatidylethanolamine, lysophosphatidylcholine in erythrocyte membranes in acute and chronic glomerulonephritis, compared with those in the control group, their information content is high - 96, 88, 21, respectively.

The study of thyroid function in patients with acute glomerulonephritis with long-term remission: the level of thyroxine-bound globulin - 0.52 ± 0.086 nmol / l, triiodothyronine 1.32 ± 0.15 nmol / l, thyroxine - 99.96 ± 7.13 nmol / l, no significant difference was found in patients

with AGN and CGN. However, if we take into account that the sum of prognostic coefficients is reliable at a value of 19.5, then a significant difference between patients of both groups is revealed.

The next stage of the study was the analysis of information content, which reflects the degree of approximation of the diagnosis to the correct diagnostic threshold. An informative measure of more than 3 was recognized as highly informative, but not less than 1, because 3 - 4 such signs are sufficient to reach the +10 threshold, i.e. providing no more than 10% errors. Such signs, for example, were a decrease in alpha - globulin, an increase in gamma - globulin, hyperlipidemia, an increase in cholesterol. Signs such as hypercoagulability, a decrease in endogenous creatinine against the background of a hereditary burden of kidney pathology, an increase in malondialdehyde and lysophosphatidylcholine, a decrease in phosphatidylethanolamine and phosphatidylcholine were prognostically unfavorable indicators. These indicators characterize the high activity of lipid peroxidation, leading to destabilization of cytomembranes and the formation of a prognostically unfavorable course of nephrotic syndrome in patients with glomerulonephritis.

Conclusions

1. In the development of chronic renal failure in children, the proportion of glomerulonephritis is still significant.
2. The peculiarity of the course of glomerulonephritis in children is that the disease develops against the background of hereditary predisposition.
3. The chronicity of the disease is due to the instability of cytomembranes against the background of increased activity of lipid peroxidation.
4. In connection with the established facts, timely diagnosis of factors leading to chronicity is necessary, as well as prevention of the formation of cytomembrane instability with immunological control of the treatment.

LITERATURE

1. Voznesenskaya T.S., Sergeeva T.V. Features of the course and risk factors for the progression of idiopathic IgA nephropathy in children. Russian Pediatric Journal. -2012.- №3,- S. 28-34.
2. Voznesenskaya T.C., Tikhomirov EE., Averyanova NS, Tsygin AN, Sergeeva TV. Mutations of NPHS1 and NPHS2 in children with congenital and infantile nephrotic syndrome // Proceedings of the VI Congress of the Russian Society of Medical Genetics. - Rostov-on-Don, 2010, - S. 39.
3. Sh.M.Ibatova, F.Kh.Mamatkulova, N.B.Abdukadirova, Kh.M.Oblokulov, F.A.Achilova. The effectiveness of the use of apricot oil in children with rickets. // Scientific and practical journal "Questions of Science and Education", Moscow, 2019, No. 27 (76), -p.40-46.
4. Mukhin N.A. Decreased glomerular filtration rate is a general population marker of poor prognosis / N.A. Mukhin // Therapeutic archive-2007. -6. -S.5-10.
5. Tsygin A.N., Timofeeva A.G., Voznesenskaya T.S., Pankratenko T.E. Nephrotic syndrome // In the book: Rational pharmacotherapy of childhood diseases. Ed. Baranova A. A., book I.-M, 2007, - S. 1024-1026.

6. Shutov A.M. Erythrocyte sedimentation rate and arterial compliance in patients with pre-dialysis chronic renal failure / A.M. Shutov, N.I. Kondratiev, V.P. Mukhorin, T.N. Ivashkina // *Nephrology: Scientific and Practical Journal*. - St. Petersburg. Aesculapius - 2003. - N1 - C. 62-66.
7. Voznesenskaya T, Tikhomirov E, Averyanova N, Tsygin A. Mutations in NPHS1 in Russian Patients with Finnish-Type Congenital Nephrotic syndrome // *Abstracts of the 15th international congress of the International Pediatric Nephrology Association -New York 2010.*-P.1921.
8. Shoira M. Ibatova Feruza Kh. Mamatkulova, Gulnoz A. Kholikova, Dilrabo Kh. Mamatkulova. Some indicators of lipid and phosphorus-calcium metabolism in children with rickets receiving conventional treatment. *International Journal of Health Sciences*, 6(S4), 2022. 3628–3638.
9. Sh.M Ibatova, F. Kh. Mamatkulova, D. Kh. Mamatkulova, N.E Ruzikulov, F.P. Abdurasulov. Study of the Clinical Features of Giambliasis in Children. *American Journal of Medicine and Medical Sciences* 2022, 12(7): 711-714.
10. Ibatova Sh. M., Mamatkulova F. Kh., Ruzikulov N.Y. The Clinical Picture of Acute Obstructive Bronchitis in Children and the Rationale for Immunomodulatory Therapy. *International Journal of Current Research and Review*. Vol 12 Issue 17. September 2020. - P.152-155.
11. Ibatova Sh. M., F. Kh. Mamatkulova, N. B. Abdukadirova, Yu. A. Rakhmonov, M. M. Kodirova. Risk Factors for Development of Broncho-Ostructive Syndrome in Children. *International Journal of Current Research and Review*. Vol 12. Issue 23 December 2020.-P. 3-6.
12. Ibatova Sh.M., Mamatkulova F.Kh., Rakhmonov Y.A., Shukurova D.B., Kodirova M.M. Assessment of the Effectiveness of Treatment of Rachit in Children by Gas-Liquid Chromatography. *International Journal of Current Research and Review*. Vol 13, Issue 06, 20 March 2021. P.64-66.
13. Sh.M. Ibatova, F.Kh. Mamatkulova, D.S. Islamova. Efficiency of combined application of apricot oil and aevit as a regulator of lipase activity of blood serum in children with vitamin D-deficiency rickets. *Journal of Critical Reviews*. // ISSN- 2394-5125. VOL 7, ISSUE 11, 2020. P.1266-1274.
14. Ibatova Sh.M., Baratova R.Sh., Mamatkulova F.Kh., Ergashev A.Kh. State of immunity in chronic obstructive pulmonary disease in children. *Asian Journal of Multidimensional Research (AJMR)*. Vol.10, Issue 3, March, 2021. P. 132-136.
15. Sh.M. Ibatova, F.Kh. Mamatkulova, N.Y. Ruzikulov, Yu.A. Rakhmonov. Bronchoo structive syndrome in children: prevalence and difficulties of differential diagnostics. *ACADEMICIA: An International Multidisciplinary Research Journal* 2021, P. 87-92.
16. Shoira M. Ibatova Feruza Kh. Mamatkulova, Gulnoz A. Kholikova, Dilrabo Kh. Mamatkulova. Some indicators of lipid and phosphorus-calcium metabolism in children with rickets receiving conventional treatment. *International Journal of Health Sciences*, 6(S4), 2022. 3628–3638.
17. Sh.M Ibatova, F. Kh. Mamatkulova, D. Kh. Mamatkulova, N.E Ruzikulov, F.P. Abdurasulov. Study of the Clinical Features of Giambliasis in Children. *American Journal of Medicine and Medical Sciences* 2022, 12(7): 711-714.